

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Lewis Y (Tumor Marker) (A70-A/A9), CF568 conjugate, 0.1mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal BOT-BNC681412-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Lewis Y (Tumor Marker) (A70-A/A9), CF568 conjugate, 0.1mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC681412-100   |
| Hersteller Artikelnummer | BNC681412-100   |
| Alternativnummer         | BOT-BNC681412-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Live Ls174T cells (human colon carcinoma cell line)   |
| Konjugation              | CF568   |
| Produktbeschreibung      | This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucalpha1-2Galbeta1-4/3GlcNAcbeta). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classification of human ren... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [A70-A/A9]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | Multiple  |
| UniProt                | Not Applicable  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |