

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

**von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)
(VWF/1465), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [VWF/1465], Mouse,
Monoclonal
BOT-BNC681465-100**

Artikelname	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1465), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [VWF/1465], Mouse, Monoclonal
Artikelnummer	BOT-BNC681465-100
Hersteller Artikelnummer	BNC681465-100
Alternativnummer	BOT-BNC681465-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human vWF protein (aa1815-1939) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[VWF/1465]
Molekulargewicht	250 kDa
Isotyp	IgG1
UniProt	P04275
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Western blotting 0.5-1 ug/mL Immunoprecipitation 0.5-1 ug/500ug protein lysate Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user