

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IgM (Immunoglobulin Mu Heavy Chain) (B-Cell Marker)(IGHM/1623), CF568 conjugate, 0.1mg/mL, Clone: [IGHM/1623], Mouse, Monoclonal BOT-BNC681623-100

| | |
|--------------------------|---|
| Artikelname | Human IgM (Immunoglobulin Mu Heavy Chain) (B-Cell Marker)(IGHM/1623), CF568 conjugate, 0.1mg/mL, Clone: [IGHM/1623], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681623-100 |
| Hersteller Artikelnummer | BNC681623-100 |
| Alternativnummer | BOT-BNC681623-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Heavy chain of human IgM |
| Konjugation | CF568 |
| Produktbeschreibung | Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric Ig... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [IGHM/1623] |

| | |
|------------------------|--|
| Molekulargewicht | 50-75 kDa |
| UniProt | P01871 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |