

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Rb1 (Tumor Suppressor Protein) (1F8), CF568 conjugate, 0.1mg/mL, Clone: [1F8], Mouse, Monoclonal BOT-BNC681701-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Rb1 (Tumor Suppressor Protein) (1F8), CF568 conjugate, 0.1mg/mL, Clone: [1F8], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC681701-100   |
| Hersteller Artikelnummer | BNC681701-100   |
| Alternativnummer         | BOT-BNC681701-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human, Mouse  |
| Immunogen                | Recombinant human Rb1 protein fragment of 283 amino acid residues (exact sequence is proprietary)   |
| Konjugation              | CF568   |
| Produktbeschreibung      | This antibody recognizes a 105 kDa phosphoprotein, identified as retinoblastoma (Rb) gene product. Its epitope is localized between aa 703-772. It shows no cross reaction with p107 or p130. It specifically stains the nuclei of BT-20 cells and primary... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |  |
|------------------------|--|
| Klon-Bezeichnung       | [1F8]  |
| Molekulargewicht       | 105 kDa  |
| UniProt                | <a href="#">P06400</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Western blotting 0.5-1 ug/mL Immunoprecipitation 0.5-1 ug/500ug protein lysate Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |