

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 6 (KRT6) (Hyperproliferation-Related Keratin) (KRT6/1702), CF568 conjugate, 0.1mg/mL, Clone: [KRT6/1702], Mouse, Monoclonal BOT-BNC681702-100**

|                          |  |
|--------------------------|--|
| Artikelname              | Cytokeratin 6 (KRT6) (Hyperproliferation-Related Keratin) (KRT6/1702), CF568 conjugate, 0.1mg/mL, Clone: [KRT6/1702], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC681702-100  |
| Hersteller Artikelnummer | BNC681702-100  |
| Alternativnummer         | BOT-BNC681702-100-100UL  |
| Hersteller               | Biotium  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | IHC  |
| Spezies Reaktivität      | Human, Mouse   |
| Immunogen                | Recombinant human full-length keratin 6 protein  |
| Konjugation              | CF568  |
| Produktbeschreibung      | This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6 (CK6). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns, and Cytokeratin 6A is the dominant... |
| Klonalität               | Monoclonal   |
| Konzentration            | 0.1 mg/mL  |

|                        |  |
|------------------------|--|
| Klon-Bezeichnung       | [KRT6/1702]  |
| Molekulargewicht       | 56 kDa   |
| UniProt                | <a href="#">P02538</a> , <a href="#">P48668</a> , <a href="#">P04259</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |