

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgM Immunoglobulin(Cocktail), CF488A conjugate, 0.1mg/mL, Clone: [Cocktail], Mouse, Monoclonal BOT-BNC880762-100

| | |
|--------------------------|---|
| Artikelname | IgM Immunoglobulin(Cocktail), CF488A conjugate, 0.1mg/mL, Clone: [Cocktail], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880762-100 |
| Hersteller Artikelnummer | BNC880762-100 |
| Alternativnummer | BOT-BNC880762-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant heavy chain of human IgM (IM260 & IM373), Heavy chain of human IgM (ICO-30) |
| Konjugation | CF488A |
| Produktbeschreibung | Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [Cocktail] |
| Molekulargewicht | 50-75 kDa |
| UniProt | P01871 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |