

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Phosphotyrosine(PY793), CF488A conjugate, 0.1mg/mL, IgG2b, Clone: [PY793], Mouse, Monoclonal BOT-BNC880793-500

| | |
|--------------------------|---|
| Artikelname | Phosphotyrosine(PY793), CF488A conjugate, 0.1mg/mL, IgG2b, Clone: [PY793], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880793-500 |
| Hersteller Artikelnummer | BNC880793-500 |
| Alternativnummer | BOT-BNC880793-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | All |
| Immunogen | Phosphotyrosine conjugated to BSA |
| Konjugation | CF488A |
| Produktbeschreibung | Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [PY793] |

| | |
|------------------------|--|
| Molekulargewicht | Depends upon the phosphorylated target |
| Isotyp | IgG2b |
| UniProt | Not Applicable |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL No pretreatment is required for staining of formalin-fixed, paraffin-embedded tissues Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |