

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD35 / CR1(CR1/802), CF488A conjugate, 0.1mg/mL, Clone: [CR1/802], Mouse, Monoclonal BOT-BNC880802-500

| | |
|--------------------------|---|
| Artikelname | CD35 / CR1(CR1/802), CF488A conjugate, 0.1mg/mL, Clone: [CR1/802], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880802-500 |
| Hersteller Artikelnummer | BNC880802-500 |
| Alternativnummer | BOT-BNC880802-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Monkey, Primate |
| Immunogen | Recombinant human CR1 protein |
| Konjugation | CF488A |
| Produktbeschreibung | Recognizes a protein of 210-220 kDa, which is identified as the complement receptor 1 (CR1)/CD35. This MAb does not block CR1 activity. It is highly specific to CR1 and shows no cross-reaction with CR2. The primary function of CR1 is to serve as the ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CR1/802] |

| | |
|------------------------|--|
| Molekulargewicht | 210-220 kDa |
| UniProt | P17927 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |