

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Caldesmon, HMW (h-Caldesmon)(CALD1/820), CF488A conjugate, 0.1mg/mL, Clone: [CALD1/820], Mouse, Monoclonal BOT-BNC880820-500

| | |
|--------------------------|---|
| Artikelname | Caldesmon, HMW (h-Caldesmon)(CALD1/820), CF488A conjugate, 0.1mg/mL, Clone: [CALD1/820], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880820-500 |
| Hersteller Artikelnummer | BNC880820-500 |
| Alternativnummer | BOT-BNC880820-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CALD1 protein |
| Konjugation | CF488A |
| Produktbeschreibung | Caldesmon HMW is the high molecular weight variant of Caldesmon. Two closely related variants of human caldesmon have been identified which are different in their electrophoretic mobility and cellular distribution. The h-caldesmon variant (120-150 kD... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CALD1/820] |

| | |
|------------------------|--|
| Molekulargewicht | 150 kDa |
| UniProt | Q05682 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |