

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD68(LAMP4/824), CF488A conjugate, 0.1mg/mL, Clone: [LAMP4/824], Mouse, Monoclonal BOT-BNC880824-100

| | |
|--------------------------|---|
| Artikelname | CD68(LAMP4/824), CF488A conjugate, 0.1mg/mL, Clone: [LAMP4/824], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880824-100 |
| Hersteller Artikelnummer | BNC880824-100 |
| Alternativnummer | BOT-BNC880824-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human LAMP4 protein |
| Konjugation | CF488A |
| Produktbeschreibung | This antibody recognizes a glycoprotein of 110 kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [LAMP4/824] |

| | |
|------------------------|--|
| Molekulargewicht | ~110 kDa |
| UniProt | P34810 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |