

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Involucrin(IVRN/827), CF488A conjugate, 0.1mg/mL, Clone: [IVRN/827], Mouse, Monoclonal BOT-BNC880827-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Involucrin(IVRN/827), CF488A conjugate, 0.1mg/mL, Clone: [IVRN/827], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC880827-500   |
| Hersteller Artikelnummer | BNC880827-500   |
| Alternativnummer         | BOT-BNC880827-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Purified involucrin from human keratinocytes  |
| Konjugation              | CF488A  |
| Produktbeschreibung      | This antibody recognizes a protein of 66 kDa-170 kDa, identified as involucrin. In Western blotting of cultured human keratinocytes, this MAb reacts with a 120 kDa protein. Involucrin is expressed in a range of stratified squamous epithelia, includin... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [IVRN/827]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 66-170 kDa  |
| UniProt                | <a href="#">P07476</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1mg/mL PBS for 5 min at 37C Flow Cytometry 0.5-1 ug/million cells/0.1 mL Note than enzyme digestion is better than citrate buffer based epitope unmasking Optimal dilution for a specific application should be determined by user |