

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p57 / KIP2(KP10), CF488A conjugate, 0.1mg/mL, Clone: [KP10], Mouse, Monoclonal BOT-BNC880879-500

| | |
|--------------------------|---|
| Artikelname | p57 / KIP2(KP10), CF488A conjugate, 0.1mg/mL, Clone: [KP10], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC880879-500 |
| Hersteller Artikelnummer | BNC880879-500 |
| Alternativnummer | BOT-BNC880879-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human p57Kip2 protein |
| Konjugation | CF488A |
| Produktbeschreibung | Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as a... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KP10] |

| | |
|------------------------|---|
| Molekulargewicht | 57 kDa |
| UniProt | P49918 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |