

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Neurofilament, phospho (NF-H) (Neuronal Marker)(NE14), CF488A conjugate, 0.1mg/mL, Clone: [NE14], Mouse, Monoclonal BOT-BNC881253-100

Artikelname	Neurofilament, phospho (NF-H) (Neuronal Marker)(NE14), CF488A conjugate, 0.1mg/mL, Clone: [NE14], Mouse, Monoclonal
Artikelnummer	BOT-BNC881253-100
Hersteller Artikelnummer	BNC881253-100
Alternativnummer	BOT-BNC881253-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Crude neurofilament preparation from porcine spinal cord
Konjugation	CF488A
Produktbeschreibung	This MAb reacts with a 200 kDa protein, identified as heavy sub-unit of neurofilaments (NF-H). It reacts specifically with the phosphorylated KSP/KEP segment at the C-terminus of the heavy subunit (NF-H) of neurofilaments. After dephosphorylation of ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[NE14]

Molekulargewicht	200 kDa
UniProt	P12036
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.25-0.5 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user