

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Prostate Specific Antigen (PSA)(KLK3/801 + KLK3/1248), CF488A conjugate, 0.1mg/mL, Clone: [KLK3/801 KLK3/1248], Mouse, Monoclonal BOT-BNC881256-500

| | |
|--------------------------|--|
| Artikelname | Prostate Specific Antigen (PSA)(KLK3/801 + KLK3/1248), CF488A conjugate, 0.1mg/mL, Clone: [KLK3/801 KLK3/1248], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC881256-500 |
| Hersteller Artikelnummer | BNC881256-500 |
| Alternativnummer | BOT-BNC881256-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KLK3 protein |
| Konjugation | CF488A |
| Produktbeschreibung | Recognizes a single protein of 33-34 kDa, identified as the prostate specific antigen (PSA). This MAbs is highly specific to PSA and stains prostatic secretory and ductal epithelium in both normal and neoplastic tissues. PSA is a chymotrypsin-like ser... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [KLK3/801 KLK3/1248] |
| Molekulargewicht | 33-34 kDa |
| UniProt | P07288 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |