

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 10 (Suprabasal Epithelial Marker) (DE-K10), CF488A conjugate, 0.1mg/mL, Clone: [DE-K10], Mouse, Monoclonal BOT-BNC881273-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 10 (Suprabasal Epithelial Marker) (DE-K10), CF488A conjugate, 0.1mg/mL, Clone: [DE-K10], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC881273-100 |
| Hersteller Artikelnummer | BNC881273-100 |
| Alternativnummer | BOT-BNC881273-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Canine, Feline, Human |
| Immunogen | Cytoskeletal preparation extracted from human ectocervical epithelium |
| Konjugation | CF488A |
| Produktbeschreibung | This MAb recognizes a protein of 56.5 kDa, identified as cytokeratin 10 (CK10). CK10 is expressed in all suprabasal layers of the epidermis. In the epidermis, expression of CK10 strictly parallels the extent of differentiation, it is absent in the ba... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [DE-K10] |
| Molekulargewicht | 56.5 kDa |
| UniProt | P13645 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |