

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), CF488A conjugate, 0.1mg/mL, Clone: [IGEL/1497R], Rabbit, Monoclonal BOT-BNC881497-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), CF488A conjugate, 0.1mg/mL, Clone: [IGEL/1497R], Rabbit, Monoclonal  |
| Artikelnummer            | BOT-BNC881497-100   |
| Hersteller Artikelnummer | BNC881497-100   |
| Alternativnummer         | BOT-BNC881497-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | FC, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human full-length MS4A1 protein   |
| Konjugation              | CF488A  |
| Produktbeschreibung      | This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD2... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [IGEL/1497R]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 33-37 kDa  |
| UniProt                | <a href="#">P11836</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |