

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HIF1 alpha (Hypoxia-Inducible Factor 1-alpha)(ESEE122), CF488A conjugate, 0.1mg/mL, Clone: [ESEE122], Mouse, Monoclonal BOT-BNC881574-100

| | |
|--------------------------|---|
| Artikelname | HIF1 alpha (Hypoxia-Inducible Factor 1-alpha)(ESEE122), CF488A conjugate, 0.1mg/mL, Clone: [ESEE122], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC881574-100 |
| Hersteller Artikelnummer | BNC881574-100 |
| Alternativnummer | BOT-BNC881574-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Bovine, Canine, Human, Mouse, Rat |
| Immunogen | GST-human HIF-1A amino acids 329-530 fusion protein |
| Konjugation | CF488A |
| Produktbeschreibung | HIF1 (hypoxia-inducible factor 1), a heterodimeric transcription factor complex central to cellular response to hypoxia, consists of two subunits (HIF-1 alpha and HIF-1 beta) which are basic helix-loop-helix proteins of the PAS (Per, ARNT, Sim) famil... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ESEE122] |

| | |
|------------------------|---|
| Molekulargewicht | 92-110 kDa |
| UniProt | Q16665 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | ELISA: For coating, purchase antibody without BSA, Flow cytometry: 0.5-1 ug/million cells, Immunofluorescence: 0.5-1 ug/mL, Optimal dilution for a specific application should be determined. Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody |