

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD8B (Cytotoxic / Suppressor T-Cell Marker)(CBB/1595), CF488A conjugate, 0.1mg/mL, Clone: [CBB/1595], Mouse, Monoclonal BOT-BNC881595-100**

Artikelname	CD8B (Cytotoxic / Suppressor T-Cell Marker)(CBB/1595), CF488A conjugate, 0.1mg/mL, Clone: [CBB/1595], Mouse, Monoclonal
Artikelnummer	BOT-BNC881595-100
Hersteller Artikelnummer	BNC881595-100
Alternativnummer	BOT-BNC881595-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Bovine, Goat, Sheep
Immunogen	Antibody was raised against bovine CD8b.
Konjugation	CF488A
Produktbeschreibung	The CD8 beta chain associates with the CD8 alpha chain to form the CD8 alpha/beta heterodimer expressed on the surface of a majority of thymocytes, and on peripheral cytotoxic alpha beta TCR T cells. This monoclonal antibody identifies cytotoxic/supp...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CBB/1595]
Molekulargewicht	32 kDa

UniProt	<a href="#">P10966</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry