

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan (Epithelial Marker) (KRTH/1576R + KRTL/1577R), CF488A conjugate, 0.1mg/mL, Clone: [KRTH/1576R KRTL/1577R], Rabbit, Monoclonal BOT-BNC881600-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin, pan (Epithelial Marker) (KRTH/1576R + KRTL/1577R), CF488A conjugate, 0.1mg/mL, Clone: [KRTH/1576R KRTL/1577R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC881600-100 |
| Hersteller Artikelnummer | BNC881600-100 |
| Alternativnummer | BOT-BNC881600-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KRT76 and KRT77 proteins |
| Konjugation | CF488A |
| Produktbeschreibung | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI <5.7) and basic (pI >6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which 67 kDa (CK1), 6... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [KRTH/1576R KRTL/1577R] |
| Molekulargewicht | 40-67 kDa |
| UniProt | Q7Z794 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution for a specific application should be determined by user |