

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Thyroglobulin(6E1.), 0.2mg/mL, Clone: [6E1.], Mouse, Monoclonal BOT-BNUB0051-500

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Thyroglobulin(6E1.), 0.2mg/mL, Clone: [6E1.], Mouse, Monoclonal                                                                                                                                                                                               |
| Artikelnummer            | BOT-BNUB0051-500                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB0051-500                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB0051-500-500UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Human thyroid follicular cells                                                                                                                                                                                                                                |
| Produktbeschreibung      | Thyroglobulin is a 660 kDa dimeric pre-protein with mutiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo c... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [6E1.]                                                                                                                                                                                                                                                        |
| Molekulargewicht         | 660 kDa (Dimeric Form)                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">P01266</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |