

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CDw17(HuLy-m13), 0.2mg/mL, IgM, Clone: [HuLy-m13], Mouse, Monoclonal BOT-BNUB1088-500

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CDw17(HuLy-m13), 0.2mg/mL, IgM, Clone: [HuLy-m13], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1088-500 |
| Hersteller Artikelnummer | BNUB1088-500 |
| Alternativnummer | BOT-BNUB1088-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | Functional Studies, IP, WB |
| Spezies Reaktivität | Human |
| Immunogen | ?-2 Microglobulin associated proteins from a detergent lysate of human PBL |
| Produktbeschreibung | CDw17 / CD17 is an intermediate glycosphingolipid from the metabolism of higher gangliosides that localizes to sphingolipid-sterol rafts. CD17 is detectable in monocytes, granulocytes, basophils, platelets, a subset of peripheral B cells (CD19) and ... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [HuLy-m13] |
| Molekulargewicht | Not Known |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgM |
| UniProt | Not Known |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |