

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

pS2(TFF1/1091), 0.2mg/mL, Clone: [TFF1/1091], Mouse, Monoclonal BOT-BNUB1091-500

| | |
|--------------------------|---|
| Artikelname | pS2(TFF1/1091), 0.2mg/mL, Clone: [TFF1/1091], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1091-500 |
| Hersteller Artikelnummer | BNUB1091-500 |
| Alternativnummer | BOT-BNUB1091-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Monkey |
| Immunogen | A synthetic peptide from the C-terminus of human TFF1 protein. |
| Produktbeschreibung | This antibody recognizes a polypeptide of 6.5 kDa, identified as pS2 estrogen-regulated protein. Its epitope is located in the c-terminus of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throu... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [TFF1/1091] |
| Molekulargewicht | 6.5 kDa |
| UniProt | P04155 |

| | |
|------------------------|--|
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |