

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), 0.2mg/mL, Clone: [IGEL/1497R], Rabbit, Monoclonal BOT-BNUB1497-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD20 / MS4A1 (B-Cell Marker) (IGEL/1497R), 0.2mg/mL, Clone:<br>[IGEL/1497R], Rabbit, Monoclonal                                                                                                                                                               |
| Artikelnummer            | BOT-BNUB1497-500                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB1497-500                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB1497-500-500UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IHC                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human full-length MS4A1 protein                                                                                                                                                                                                                   |
| Produktbeschreibung      | This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD2... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [IGEL/1497R]                                                                                                                                                                                                                                                  |
| Molekulargewicht         | 33-37 kDa                                                                                                                                                                                                                                                     |

|                        |                                                                                                                                                                                                                                                                                                              |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">P11836</a>                                                                                                                                                                                                                                                                                       |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                  |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                       |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |