

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD1a / HTA1 (Mature Langerhans Cells Marker) (C1A/1506R), 0.2mg/mL, Clone: [C1A/1506R], Rabbit, Monoclonal BOT-BNUB1506-500

Artikelname	CD1a / HTA1 (Mature Langerhans Cells Marker) (C1A/1506R), 0.2mg/mL, Clone: [C1A/1506R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB1506-500
Hersteller Artikelnummer	BNUB1506-500
Alternativnummer	BOT-BNUB1506-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length CD1a protein
Produktbeschreibung	At least five CD1 genes (CD1a, b, c, d, and e) are identified. CD1 proteins have been demonstrated to restrict T cell response to non-peptide lipid and glycolipid antigens and play a role in non-classical antigen presentation. CD1a is a non-polymorph...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[C1A/1506R]
Molekulargewicht	49 kDa

UniProt	P06126
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user