

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Catenin, beta (p120) (CTNNB1/1508), 0.2mg/mL, Clone: [CTNNB1/1508], Mouse, Monoclonal BOT-BNUB1508-500

Artikelname	Catenin, beta (p120) (CTNNB1/1508), 0.2mg/mL, Clone: [CTNNB1/1508], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1508-500
Hersteller Artikelnummer	BNUB1508-500
Alternativnummer	BOT-BNUB1508-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant human full-length beta-Catenin (p120) protein fragment
Produktbeschreibung	Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[CTNNB1/1508]

Molekulargewicht	92 kDa
UniProt	P35222
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user