

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD45 / LCA (B-Cell Marker) (F10-89-4), 0.2mg/mL, Clone: [F10-89-4], Mouse, Monoclonal BOT-BNUB1522-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD45 / LCA (B-Cell Marker) (F10-89-4), 0.2mg/mL, Clone: [F10-89-4],<br>Mouse, Monoclonal                                                                                                                                                                      |
| Artikelnummer            | BOT-BNUB1522-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB1522-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB1522-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IF, IHC                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Human T-Lymphocytes                                                                                                                                                                                                                                           |
| Produktbeschreibung      | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [F10-89-4]                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 180-220 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| UniProt                | <a href="#">P08575</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |