

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NKX2.2 (Neuroendocrine and Ewings Sarcoma Marker) (NX2/1524), 0.2mg/mL, Clone: [NX2/1524], Mouse, Monoclonal BOT-BNUB1524-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | NKX2.2 (Neuroendocrine and Ewings Sarcoma Marker) (NX2/1524), 0.2mg/mL, Clone: [NX2/1524], Mouse, Monoclonal                                                                                                                                                  |
| Artikelnummer            | BOT-BNUB1524-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB1524-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB1524-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Human recombinant NKX2.2 protein fragment (aa1-119) (exact sequence is proprietary)                                                                                                                                                                           |
| Produktbeschreibung      | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [NX2/1524]                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 40-50 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| UniProt                | <a href="#">O95096</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |