

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100B (Astrocyte and Melanoma Marker) (S100B/1706R), 0.2mg/mL, Clone: [S100B/1706R], Rabbit, Monoclonal BOT-BNUB1706-500

Artikelname	S100B (Astrocyte and Melanoma Marker) (S100B/1706R), 0.2mg/mL, Clone: [S100B/1706R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB1706-500
Hersteller Artikelnummer	BNUB1706-500
Alternativnummer	BOT-BNUB1706-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	Recombinant full-length human S100B protein
Produktbeschreibung	S100A and S100B proteins are two members of the S100 family of calcium binding proteins. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the beta-cha...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[S100B/1706R]
Molekulargewicht	10-12 kDa

UniProt	P04271
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user