

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8 (KRT8) (KRT8/2174R), 0.2mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal BOT-BNUB2174-500**

Artikelname	Cytokeratin 8 (KRT8) (KRT8/2174R), 0.2mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2174-500
Hersteller Artikelnummer	BNUB2174-500
Alternativnummer	BOT-BNUB2174-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Porcine, Rabbit, Sheep
Immunogen	Recombinant full-length human KRT8 protein
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[KRT8/2174R]
Molekulargewicht	52.5 kDa

UniProt	<a href="#">P05787</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user