

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) (rGROEL/780), 0.2mg/mL, Clone: [rGROEL/780], Mouse, Monoclonal BOT-BNUB2207-500

| | |
|--------------------------|---|
| Artikelname | HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) (rGROEL/780), 0.2mg/mL, Clone: [rGROEL/780], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB2207-500 |
| Hersteller Artikelnummer | BNUB2207-500 |
| Alternativnummer | BOT-BNUB2207-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Bovine, Canine, Gallus, Hamster, Human, Invertebrate, Monkey, Mouse, Porcine, Rabbit, Rat, Sheep, Xenopus |
| Immunogen | Recombinant full-length human HSP60 protein |
| Produktbeschreibung | This antibody recognizes a 60 kDa protein, identified as the heat shock protein 60 (hsp60). Its epitope is localized between aa 383-447 of human hsp60. A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis ... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [rGROEL/780] |

| | |
|------------------------|---|
| Molekulargewicht | 60 kDa |
| UniProt | P10809 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.25-0.5 ug/mL Optimal dilution for a specific application should be determined by user |