

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 0.2mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal BOT-BNUB2748-100

Artikelname	AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 0.2mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2748-100
Hersteller Artikelnummer	BNUB2748-100
Alternativnummer	BOT-BNUB2748-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human AMACR protein fragment (around aa 297-394) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 42 kDa, which is identified as Alpha-methylacyl-CoA Racemase (AMACR), also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and betbeta-oxidation of branched-chain fatty acids. AMACR is ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[AMACR/2748R]

Molekulargewicht	42 kDa
UniProt	Q9UHK6
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user