

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100A4 (Marker of Tumor Metastasis) (S100A4/2750R), 0.2mg/mL, Clone: [S100A4/2750R], Rabbit, Monoclonal BOT-BNUB2750-500

Artikelname	S100A4 (Marker of Tumor Metastasis) (S100A4/2750R), 0.2mg/mL, Clone: [S100A4/2750R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2750-500
Hersteller Artikelnummer	BNUB2750-500
Alternativnummer	BOT-BNUB2750-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant fragment of human S100A4 protein (around aa1-200) (exact sequence is proprietary)
Produktbeschreibung	S100A4 belongs to the S100 super-family of proteins containing 2 EF-hand calcium-binding domains. S100 genes include at least 25 members, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z. S100A4 exerts its function via dir...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[S100A4/2750R]

Molekulargewicht	10-12 kDa
UniProt	P26447
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user