

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MAGE A1(MA454), 1mg/mL, Clone: [MA454], Mouse, Monoclonal BOT-BNUM0008-50**

|                          |   |
|--------------------------|---|
| Artikelname              | MAGE A1(MA454), 1mg/mL, Clone: [MA454], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUM0008-50   |
| Hersteller Artikelnummer | BNUM0008-50   |
| Alternativnummer         | BOT-BNUM0008-50-50UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, FC, IHC, WB  |
| Spezies Reaktivität      | Canine, Human, Rat  |
| Immunogen                | Human MAGE-A1 full length recombinant protein   |
| Produktbeschreibung      | Recognizes a protein of 42-46 kDa, identified as MAGE-1. This MAb does not cross-react with MAGE-2, -3, -4, -6 -9, -10, -or -12 protein. Human malignant neoplasms carry rejection antigens that are recognized by the patients autologous, tumor directed... |
| Klonalität               | Monoclonal  |
| Konzentration            | 1 mg/mL   |
| Klon-Bezeichnung         | [MA454]   |
| Molekulargewicht         | 42-46 kDa   |
| UniProt                  | <a href="#">P43355</a>  |

|                        |  |
|------------------------|--|
| Puffer                 | PBS, no BSA, no azide  |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |