

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MART-1 / Melan-A(M2-9E3), 1mg/mL, Clone: [M2-9E3], Mouse, Monoclonal BOT-BNUM0010-50

| | |
|--------------------------|---|
| Artikelname | MART-1 / Melan-A(M2-9E3), 1mg/mL, Clone: [M2-9E3], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0010-50 |
| Hersteller Artikelnummer | BNUM0010-50 |
| Alternativnummer | BOT-BNUM0010-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant hMART-1 protein |
| Produktbeschreibung | This antibody recognizes a protein doublet of 20-22 kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Sev... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [M2-9E3] |
| Molekulargewicht | 20-22 kDa (doublet) |

| | |
|------------------------|---|
| UniProt | Q16655 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |