

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 18(DC10), 1mg/mL, IgG1, Clone: [DC10], Mouse, Monoclonal BOT-BNUM0021-50

Artikelname	Cytokeratin 18(DC10), 1mg/mL, IgG1, Clone: [DC10], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0021-50
Hersteller Artikelnummer	BNUM0021-50
Alternativnummer	BOT-BNUM0021-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	PMC-42 human breast carcinoma cells
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[DC10]
Molekulargewicht	45 kDa

Isotyp	IgG1
UniProt	P05783
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody conjugates</p> <p>Immunofluorescence: 1-2 ug/mL</p> <p>Immunohistology formalin-fixed 0.5-1 ug/mL</p> <p>Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes</p> <p>Flow Cytometry 0.5-1 ug/million cells/0.1 mL</p> <p>Western blotting 0.5-1 ug/mL</p> <p>Does not react with mouse, rat, ferret, hamster, sheep, cow, dog or pig, others not known</p> <p>Optimal dilution for a specific application should be determined by user</p>