

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD35 / CR1(E11), 1mg/mL, Clone: [E11], Mouse, Monoclonal BOT-BNUM0040-50

| | |
|--------------------------|---|
| Artikelname | CD35 / CR1(E11), 1mg/mL, Clone: [E11], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0040-50 |
| Hersteller Artikelnummer | BNUM0040-50 |
| Alternativnummer | BOT-BNUM0040-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IHC, WB |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Intact human monocytes |
| Produktbeschreibung | Recognizes a protein of 210-220 kDa, which is identified as the complement receptor 1 (CR1)/CD35. This MAb is specific for a site in CR1 that is distal from the C3b-binding site, so that it is unable to block CR1 activity. This MAb is highly specific... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [E11] |
| Molekulargewicht | 210-220 kDa |
| UniProt | P17927 |

| | |
|------------------------|---|
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |