

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Granulocyte Marker(BM-2), 1mg/mL, IgG1, Clone: [BM-2], Mouse, Monoclonal BOT-BNUM0062-50**

Artikelname	Granulocyte Marker(BM-2), 1mg/mL, IgG1, Clone: [BM-2], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0062-50
Hersteller Artikelnummer	BNUM0062-50
Alternativnummer	BOT-BNUM0062-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes
Produktbeschreibung	Recognizes 183 kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BM-2]

Molekulargewicht	183 kDa
Isotyp	IgG1
UniProt	Not Known
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>