

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD57 / B3GAT1(NK-1 + HNK-1), 1mg/mL, IgM, Clone: [NK-1 HNK-1], Mouse, Monoclonal BOT-BNUM0695-50

Artikelname	CD57 / B3GAT1(NK-1 + HNK-1), 1mg/mL, IgM, Clone: [NK-1 HNK-1], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0695-50
Hersteller Artikelnummer	BNUM0695-50
Alternativnummer	BOT-BNUM0695-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human peripheral blood mononuclear cells (HNK-1 & NK-1)
Produktbeschreibung	Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[NK-1 HNK-1]
Molekulargewicht	~110 kDa

Isotyp	IgM
UniProt	Q9P2W7
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user