

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TNF alpha(TNF706), 1mg/mL, Clone: [TNF706], Mouse, Monoclonal BOT-BNUM0706-50

| | |
|--------------------------|---|
| Artikelname | TNF alpha(TNF706), 1mg/mL, Clone: [TNF706], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM0706-50 |
| Hersteller Artikelnummer | BNUM0706-50 |
| Alternativnummer | BOT-BNUM0706-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Canine, Feline, Human, Mouse, Rabbit, Rat, Zebrafish |
| Immunogen | Recombinant N-terminal fragment of human TNF-? |
| Produktbeschreibung | Tumor Necrosis Factor Alpha (TNF alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha is believed to mediate pathogenic shock and tissue injury associate... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [TNF706] |
| Molekulargewicht | 17 kDa |
| UniProt | P01375 |

| | |
|------------------------|--|
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-paraffin 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |