

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53(BP53-12 + DO-7), 1mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal BOT-BNUM0723-50

Artikelname	p53(BP53-12 + DO-7), 1mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0723-50
Hersteller Artikelnummer	BNUM0723-50
Alternativnummer	BOT-BNUM0723-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Bovine, Human, Monkey
Immunogen	Recombinant human wild-type p53 protein (BP53-12) & Recombinant human wild type p53 protein expressed in E. coli (DO-7)
Produktbeschreibung	Recognizes a 53 kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53 under denaturing and non-denaturing conditions. Its epitope maps within the N-terminus (aa 20-25) of p53 oncopr...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BP53-12 DO-7]
Molekulargewicht	53 kDa

UniProt	P04637
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user