

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

bcl-x(2H12), 1mg/mL, IgG2a, Clone: [2H12], Mouse, Monoclonal BOT-BNUM0751-50

Artikelname	bcl-x(2H12), 1mg/mL, IgG2a, Clone: [2H12], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0751-50
Hersteller Artikelnummer	BNUM0751-50
Alternativnummer	BOT-BNUM0751-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Mouse, Porcine, Rat
Immunogen	A synthetic peptide, aa 3-14 (Cys-QSNRELVVDFLS) of human Bcl-X protein.
Produktbeschreibung	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[2H12]
Molekulargewicht	27 kDa

Isotyp	IgG2a
UniProt	Q07817
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user