

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27 / KIP1(DCS-72.F6), 1mg/mL, Clone: [DCS-72.F6], Mouse, Monoclonal BOT-BNUM0771-50

Artikelname	p27 / KIP1(DCS-72.F6), 1mg/mL, Clone: [DCS-72.F6], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0771-50
Hersteller Artikelnummer	BNUM0771-50
Alternativnummer	BOT-BNUM0771-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Human, Monkey, Mouse, Rat
Immunogen	Mouse recombinant p27 protein
Produktbeschreibung	Recognizes a 27 kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. Its epitope spans between aa 83-204 of p27. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27Kip1 functions ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[DCS-72.F6]
Molekulargewicht	25-26 kDa

UniProt	P46527
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user