

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p34 / cdk1(POH-1), 1mg/mL, Clone: [POH-1], Mouse, Monoclonal BOT-BNUM0853-50

Artikelname	p34 / cdk1(POH-1), 1mg/mL, Clone: [POH-1], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0853-50
Hersteller Artikelnummer	BNUM0853-50
Alternativnummer	BOT-BNUM0853-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mink, Monkey
Immunogen	Purified recombinant human p34cdc2 fusion protein
Produktbeschreibung	Recognizes a 34 kDa protein, which is identified as cyclin dependent kinase 1 (cdk1) or p34cdc2 protein kinase. cdk1 / p34cdc2 plays a crucial role during cell division and is most active during mitosis. It is predominantly localized in the nucleus. ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[POH-1]
Molekulargewicht	34 kDa
UniProt	P06493

Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse, rat, Xenopus, Drosophila, S. pombe, or S. cerevisiae, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user