

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

bcl-2(BCL2/782 + BCL2/796), 1mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal BOT-BNUM1023-50

Artikelname	bcl-2(BCL2/782 + BCL2/796), 1mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1023-50
Hersteller Artikelnummer	BNUM1023-50
Alternativnummer	BOT-BNUM1023-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human bcl-2 protein
Produktbeschreibung	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BCL2/782 BCL2/796]
Molekulargewicht	25-26 kDa

UniProt	P10415
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user