

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Blood Group A(HE-10), 1mg/mL, Clone: [HE-10], Mouse, Monoclonal BOT-BNUM1089-50

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Blood Group A(HE-10), 1mg/mL, Clone: [HE-10], Mouse, Monoclonal                                                                                                                                                                                               |
| Artikelnummer            | BOT-BNUM1089-50                                                                                                                                                                                                                                               |
| Hersteller Artikelnummer | BNUM1089-50                                                                                                                                                                                                                                                   |
| Alternativnummer         | BOT-BNUM1089-50-50UL                                                                                                                                                                                                                                          |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Mixture of erythrocytes of blood group A and glycoprotein fraction isolated from the saliva of secretors with blood group A                                                                                                                                   |
| Produktbeschreibung      | This antibody is applicable for staining ABO blood groups A and AB. The histo-blood group ABO involves three carbohydrate antigens: A, B, and H. Blood group antigens are generally defined as molecules formed by sequential addition of saccharides to t... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 1 mg/mL                                                                                                                                                                                                                                                       |
| Klon-Bezeichnung         | [HE-10]                                                                                                                                                                                                                                                       |
| Molekulargewicht         | Multiple                                                                                                                                                                                                                                                      |

|                        |                                                                                                                                                                                                                                                                                                 |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">P16442</a>                                                                                                                                                                                                                                                                          |
| Puffer                 | PBS, no BSA, no azide                                                                                                                                                                                                                                                                           |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                          |
| Anwendungsbeschreibung | Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |