

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611), 1mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal BOT-BNUM1611-50

Artikelname	CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611), 1mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1611-50
Hersteller Artikelnummer	BNUM1611-50
Alternativnummer	BOT-BNUM1611-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CD22 protein (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[LPFS2/1611]
Molekulargewicht	120 kDa

Isotyp	IgG1
UniProt	P26842
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user