

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), 1mg/mL, Clone: [VWF/1767], Mouse, Monoclonal BOT-BNUM1767-50**

Artikelname	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), 1mg/mL, Clone: [VWF/1767], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1767-50
Hersteller Artikelnummer	BNUM1767-50
Alternativnummer	BOT-BNUM1767-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human vWF protein (aa1815-1939) (exact sequence is proprietary)
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[VWF/1767]

Molekulargewicht	250 kDa
UniProt	<a href="#">P04275</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user