

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD8A (Cytotoxic- & Suppressor T-Cell Marker) (C8/1779R), 1mg/mL, Clone: [C8/1779R], Rabbit, Monoclonal BOT-BNUM1779-50**

Artikelname	CD8A (Cytotoxic- & Suppressor T-Cell Marker) (C8/1779R), 1mg/mL, Clone: [C8/1779R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1779-50
Hersteller Artikelnummer	BNUM1779-50
Alternativnummer	BOT-BNUM1779-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD8a protein
Produktbeschreibung	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[C8/1779R]
Molekulargewicht	32 kDa

UniProt	<a href="#">P01732</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user