

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Histone H1 (Nuclear Marker) (HH1/1784R), 1mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal BOT-BNUM1784-50**

Artikelname	Histone H1 (Nuclear Marker) (HH1/1784R), 1mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1784-50
Hersteller Artikelnummer	BNUM1784-50
Alternativnummer	BOT-BNUM1784-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human Histone H1 protein
Produktbeschreibung	Eukaryotic histones are basic and water-soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed superhelical turn sequentially to form chromosomal fiber. Two molecules of each of...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[HH1/1784R]
Molekulargewicht	~30 kDa

UniProt	Multiple
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user