

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Epithelial Marker) (rMOC-31), 1mg/mL, Clone: [rMOC-31], Mouse, Monoclonal BOT-BNUM1819-50

| | |
|--------------------------|---|
| Artikelname | Ep-CAM / CD326 (Epithelial Marker) (rMOC-31), 1mg/mL, Clone: [rMOC-31], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1819-50 |
| Hersteller Artikelnummer | BNUM1819-50 |
| Alternativnummer | BOT-BNUM1819-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Neuraminidase treated GLS-1 human small cell lung carcinoma cells |
| Produktbeschreibung | The binding epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epith... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [rMOC-31] |
| Molekulargewicht | 40-43 kDa |

| | |
|------------------------|--|
| UniProt | P16422 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |